

ON THE SPECTRUM OF ASYMPTOTIC ENTROPIES OF RANDOM WALKS

OMER TAMUZ AND TIANYI ZHENG

ABSTRACT. Given a random walk on a free group, we study the random walks it induces on the group's quotients. We show that the spectrum of asymptotic entropies of the induced random walks has no isolated points, except perhaps its maximum.

1. INTRODUCTION

Let G be a finitely generated group, and let μ be a probability measure on G . The μ -random walk on G is a time homogeneous Markov chain g_1, g_2, \dots on the state space G whose steps are distributed i.i.d. μ : for $g, h \in G$ the transition probability from g to h is $\mu(g^{-1}h)$. An important statistic of a random walk is its *Avez Asymptotic Entropy* [3]

$$h(G, \mu) := \lim_{n \rightarrow \infty} \frac{1}{n} H(g_n),$$

where $H(\cdot)$ is the Shannon entropy. The importance of asymptotic entropy is due to the fact that it vanishes if and only if every bounded μ -harmonic function is constant; that is, if the μ -random walk has a trivial Poisson boundary [3, 29]. Moreover, as the asymptotic entropy is the limit of the mutual information $I(g_1; g_n)$ between the first step of the random walk and its position in later time periods, it quantifies the extent by which the random walk fails to have the Liouville property.

Suppose that G has d generators, and let μ be the symmetric measure that assigns $1/(2d)$ to each generator and its inverse. The main question that we ask in this paper is: what possible values of $h(G, \mu)$ are attained as we vary the group G ?

To formalize and generalize this question, we consider the following setting. Given G and μ , and given a quotient $\Gamma = G/N$, the induced random walk g_1N, g_2N, \dots on Γ has step distribution μ_Γ , where, for $\gamma = gN$, $\mu_\Gamma(\gamma) = \mu(gN)$. In other words, μ_Γ is the push-forward of μ under the quotient map; we will simply write μ instead of μ_Γ whenever this is unambiguous. For a given G and μ , what values can be realized as the asymptotic random walk entropies of such quotients? This is particularly interesting when G has many quotients, and we indeed focus on the case that of \mathbf{F}_d , the free groups with $d \geq 2$ generators.

Omer Tamuz was supported by a grant from the Simons Foundation (#419427).

Given (G, μ) we denote the spectrum of random walk entropies by

$$\mathfrak{H}(G, \mu) := \{h(\Gamma, \mu) : \Gamma \text{ is a quotient group of } G\}.$$

We will consider measures μ on G that have *finite first moment*, that is, $\sum_{g \in G} |g|_S \mu(g) < \infty$, where $|\cdot|_S$ is the word length with respect to generating set S . Recall that μ is *non-degenerate* if its support generates G as a semigroup.

Our main result is the following.

Theorem 1.1. *Let μ be a non-degenerate probability measure with finite first moment on the free group \mathbf{F}_d , $d \geq 2$. Suppose Γ is a proper quotient of \mathbf{F}_d . Then for any $\epsilon > 0$, there exists a quotient group $\tilde{\Gamma}$ of \mathbf{F}_d such that $\mathbf{F}_d \twoheadrightarrow \tilde{\Gamma} \twoheadrightarrow \Gamma$ and*

$$h(\Gamma, \mu) < h(\tilde{\Gamma}, \mu) < h(\Gamma, \mu) + \epsilon.$$

In particular, the set $\mathfrak{H}(\mathbf{F}_d, \mu)$ has no isolated points, except perhaps its maximum.

It follows from Theorem 1.1 that if $\mathfrak{H}(\mathbf{F}_d, \mu)$ is a closed subset in \mathbb{R} , then it must be the full interval $[0, h(\mathbf{F}_d, \mu)]$. To the best of our knowledge, it is not known whether the set $\mathfrak{H}(\mathbf{F}_d, \mu)$ is closed.

The key ingredient in the proof of Theorem 1.1 is an explicit construction, which might be of independent interest, of a sequence of groups in the space \mathcal{G}_d of d -marked groups with the following properties.

Proposition 1.2. *Let μ be a non-degenerate probability measure on \mathbf{F}_d , $d \geq 2$, with finite first moment. Then there exists a sequence of marked groups $((\Gamma_n, S_n))_{n=1}^\infty$ in \mathcal{G}_d such that:*

- (i): *The sequence (Γ_n, S_n) converges to $(\mathbf{F}_d, \mathbf{S})$ as $n \rightarrow \infty$ in the space of d -marked groups.*
- (ii): *The sequence of asymptotic entropies $h(\Gamma_n, \mu) \rightarrow 0$ as $k \rightarrow \infty$.*
- (iii): *For each $n \in \mathbb{N}$, Γ_n is non-amenable, has no nontrivial amenable normal subgroups, and has only countably many amenable subgroups.*

The moment condition on μ is used to bound the asymptotic entropy. It seems to be an interesting question whether Proposition 1.2 remains true assuming only that μ has finite entropy.

Property (iii) in the statement above implies that the action of Γ_n on the Poisson boundary of (Γ_n, μ) is essentially free. This property is crucial for our purposes. Any sequence of d -marked finite groups with girth growing to infinity would satisfy properties (i) and (ii), but the Poisson boundaries are trivial for finite groups.

We construct the sequence of marked groups as stated via taking extensions of the Fabrykowski-Gupta group. Necessary terminology and background are reviewed in Section 2. Provided the sequence of marked groups stated in Proposition 1.2, the proof of Theorem 1.1 is completed by taking suitable diagonal product of groups; see Section 4.

1.1. Boundary entropies. A closely related question—and, to our knowledge, a much better studied one—is that of the spectrum of *Furstenberg entropies*. Let (X, ν) be a standard Borel space, equipped with a probability measure, and on which G acts by measure class preserving transformations. We say that (X, ν) is a (G, μ) -space, if the measure ν is μ -stationary, that is $\mu * \nu = \nu$. The Furstenberg entropy of a (G, μ) -space (X, ν) is a numerical invariant defined in [18] as

$$h_\mu(X, \nu) := \sum_{g \in G} \mu(g) \int_X -\log \frac{dg^{-1}\nu}{d\nu} d\nu.$$

The *Furstenberg entropy realization problem* asks given (G, μ) , what is the spectrum of the Furstenberg entropy $h_\mu(X, \nu)$, as (X, ν) varies over all ergodic μ -stationary actions of G .

We briefly summarize what is known about this problem. In Kaimonovich and Vershik [29] it is shown that $h_\mu(X, \nu) \leq h(G, \mu)$. The Poisson boundary of an induced random walk on a quotient group G/N is a (G, μ) -space, whose Furstenberg entropy is equal to the random walk's asymptotic entropy. Hence every realizable random walk entropy value is also a realizable Furstenberg entropy value.

Nevo [36] shows that whenever G has Kazhdan's property (T) then there is a constant $c > 0$, depending on (G, μ) , such that whenever $h_\mu(X, \nu) < c$ then it in fact vanishes. In [12], Bowen showed that for the free group \mathbf{F}_d , $d \geq 2$, and μ uniform on the symmetric free generating set $\mathbf{S} \cup \mathbf{S}^{-1}$, all values in $[0, h(\mathbf{F}_d, \mu)]$ can be realized as the Furstenberg entropy of an ergodic stationary action of \mathbf{F}_d .¹

A particularly important class of (G, μ) -space are the (G, μ) -boundaries. These are the G -factors of the Poisson boundary of (G, μ) , and include the Poisson boundaries of the induced random walks on quotient groups. For such boundaries, the next result is an analogue of Theorem 1.1.

Theorem 1.3. *In the setting of Theorem 1.1, suppose (X, ν) is a (\mathbf{F}_d, μ) -boundary such that the action of \mathbf{F}_d is not essentially free. Then for any $\epsilon > 0$, there exists a (\mathbf{F}_d, μ) -boundary $(\tilde{X}, \tilde{\nu})$ such that*

$$h(X, \nu) < h(\tilde{X}, \tilde{\nu}) < h(X, \nu) + \epsilon,$$

and (X, ν) is an \mathbf{F}_d -factor of $(\tilde{X}, \tilde{\nu})$.

¹The approach in [12] is to take an ergodic invariant random subgroup of G and construct an ergodic stationary system (which can be referred to as a Poisson bundle, using the terminology introduced in [28]). The Furstenberg entropy of this stationary system is then studied by considering random walk entropies on the coset spaces associated with the invariant random subgroups. Recall that an IRS is a Borel probability measure η on the Chabauty space $\text{Sub}(G)$ of closed subgroups of G , which is invariant under conjugation by G . For further work on the Furstenberg entropy realization problem using the IRS-Poisson bundle approach, see [25, 26] and references therein.

Note that if an ergodic invariant random subgroup is not almost surely a normal subgroup, then the corresponding Poisson bundle is not a quotient of the Poisson boundary of (G, μ) because of the measure-preserving factor to the invariant random subgroup. Hence Bowen's results do not resolve the question for Furstenberg entropies of (\mathbf{F}_d, μ) -boundaries, or for asymptotic random walk entropies.

1.2. Spectral radii. The same kind of construction as in the proof of Theorem 1.1 implies the following result on spectral radii of symmetric random walks. Recall that the spectral radius of a μ -random walk on Γ is defined as

$$\rho(\Gamma, \mu) = \limsup_{2n \rightarrow \infty} \mu^{(n)}(id_\Gamma)^{\frac{1}{2n}},$$

where $\mu^{(n)}$ is the n -fold convolution of μ with itself.

Theorem 1.4. *Let μ be a symmetric non-degenerate probability measure on the free group \mathbf{F}_d , $d \geq 2$. Suppose Γ is a proper quotient of \mathbf{F}_d . Then for any $\epsilon > 0$, there exists a quotient group $\tilde{\Gamma}$ of \mathbf{F}_d such that $\mathbf{F}_d \rightarrow \tilde{\Gamma} \rightarrow \Gamma$ and*

$$\rho(\Gamma, \mu) - \epsilon < \rho(\tilde{\Gamma}, \mu) < \rho(\Gamma, \mu).$$

Our construction uses a diagonal product of marked groups, and is similar to the construction in [30]. A result of Kassabov and Pak [31] states that the set of the spectral radii $\{\rho(\Gamma, \mu) : \Gamma \text{ is a quotient of } \mathbf{F}_d\}$ contains a subset homeomorphic to the Cantor set. The same construction shows that the set $\mathfrak{H}(\mathbf{F}_d, \mu)$ contains a subset homeomorphic to the Cantor set as well. It is not known whether this set of spectral radii is closed.

Acknowledgement. We thank Michael Björklund, Jérémie Brieussel, Yair Hartman and Igor Pak for helpful discussions.

2. PRELIMINARIES

2.1. (G, μ) -boundaries. In this paper we only consider countable groups. A probability measure μ on G is non-degenerate if the support of μ generates G as a semigroup. For a countable group G , we say a Lesbesgue space (X, ν) is a G -space, if G acts measurably on X and the probability measure ν is quasi-invariant with respect to the G -action. A G -space (X, ν) is ergodic if every G -invariant subset is either null or conull. A measurable map $\pi : (X, \nu) \rightarrow (Y, \eta)$ is called a G -map if it is G -equivariant and η is the pushforward of ν under π .

Given a probability measure μ on G , let $\Omega = G^{\mathbb{N}}$ be the path space, \mathbb{P}_μ be the law of the μ -random walk starting at id , and \mathcal{I} be the σ -field on Ω that is invariant under time shifts. The Poisson boundary of (G, μ) is denoted by the measure space (B, \mathcal{F}, ν_B) together with a G -map $\mathbf{b} : (\Omega, \mathcal{I}, \mathbb{P}_\mu) \rightarrow (B, \mathcal{F}, \nu_B)$, where $\mathbf{b}^{-1}\mathcal{F} = \mathcal{I}$ up to null sets with respect to \mathbb{P}_μ , and the σ -algebra \mathcal{F} is countably generated and separating points. The existence and uniqueness up to isomorphism of the Poisson boundary of (G, μ) was shown by Furstenberg

[18, 19, 20]. The G -action on the Poisson boundary (B, ν_B) is ergodic, and in fact doubly ergodic, by Kaimanovich [27].

We use the notation (B, ν_B) to denote a compact model of the Poisson boundary of (G, μ) , which exists by the Mackey realization [34]. A (G, μ) -boundary (X, ν) is defined to be a G -factor of (B, ν_B) . Moreover, the factor map $(B, \nu_B) \rightarrow (X, \nu)$ is essentially unique, see [4, Theorem 2.14], and we will denote it by β_X .

Denote by $P(X)$ the space of Borel probability measures on the compact space X . A factor map $\pi : (Y, \eta) \rightarrow (X, \nu)$ gives a unique disintegration map $D_\pi : X \rightarrow P(Y)$ such that for ν -a.e. $x \in X$, $D_\pi(x)$ is supported on the fiber of x and $\int_X D_\pi(x) d\nu(x) = \eta$. We say (Y, η) is a *relatively measure preserving extension* of X if D_π is G -equivariant, that is $D_\pi(g \cdot x) = g \cdot D_\pi(x)$.

We will need the following properties regarding Furstenberg entropy and relatively measure preserving extensions.

Proposition 2.1 ([37, Proposition 1.9]). *Let $\pi : (Y, \eta) \rightarrow (X, \nu)$ be a G -factor map. Suppose $h(X, \nu) < \infty$ and $h(Y, \eta) = h(X, \nu)$. Then (Y, η) is a relative measure preserving extension of (X, ν) .*

Lemma 2.2 ([4, Corollary 2.20]). *Let $\pi : (Y, \eta) \rightarrow (X, \nu)$ be a relatively measure-preserving extension of two (G, μ) -boundaries. Then $(Y, \eta) = (X, \nu)$.*

2.2. The space of marked groups and convergence to the free group.

Denote by \mathcal{G}_d the space of d -generated groups (G, S) , where $S = (s_1, \dots, s_d)$ is a generating tuple, equipped with the *Cayley-Grigorchuk topology*. We refer to the pair (G, S) as a marked group and \mathcal{G}_d the space of d -marked groups. Recall that in this topology, two marked groups (G_1, S_1) and (G_2, S_2) are close if marked balls of large radius in the Cayley graphs of (G_1, S_1) and (G_2, S_2) around the identities are isomorphic. This space is introduced by Grigorchuk in [23].

Denote by $(\mathbf{F}_d, \mathbf{S})$ a free group of rank d , where $\mathbf{S} = (\mathbf{s}_1, \dots, \mathbf{s}_d)$ consists of the free generators. Let G be a d -generated group. Following the definition in Akhmedov [2] and Olshanskii-Sapir [38], we say a non-trivial word $w(x_1, \dots, x_d)$ is a *d -almost-identity* for G , if the identity $w(g_1, \dots, g_d) = 1$ is satisfied for any d -generating tuple (g_1, \dots, g_d) . By [38, Theorem 9], there exists a sequence of d -markings $(G, S_k)_{k=1}^\infty$ that converges to $(\mathbf{F}_d, \mathbf{S})$ in the Cayley-Grigorchuk topology if G is d -generated and satisfies no d -almost identity.

In [1], Abért gives a general criterion for a group to satisfy no identity. Suppose $G \curvearrowright X$ by permutations. We say G *separates* X , if for every finite subset Y of X , the pointwise fixator $G_Y = \{g \in G : y \cdot g = y \text{ for all } y \in Y\}$ has no fixed point outside Y . Abért shows that if G separates X then G satisfies no identity. Bartholdi and Erschler [6] provide a criterion for absence of almost-identities: under the additional assumption that the Frattini subgroup $\Phi(G)$ has finite index in G , the condition in Abért's criterion implies that G satisfies no almost-identity. Recall that the Frattini subgroup of G is the intersection of all the maximal subgroups of G .

Weakly branch groups provide examples of groups satisfying Abért's criterion. The notion of weakly branch group is introduced by Grigorchuk in [24]. Let \mathbb{T} be a rooted spherical symmetric tree. For a vertex $u \in \mathbb{T}$, let C_u be the set of infinite rays with prefix u . We say a group G acting by automorphisms on \mathbb{T} is *weakly branching* if it acts level transitively and the rigid stabilizer $\text{Rist}_G(C_u)$ of any vertex $u \in \mathbb{T}$ is nontrivial. Recall that $\text{Rist}_G(C_u) = \{g \in G : x \cdot g = x \text{ for all } x \notin C_u\}$, that is, the set of group elements that only move the descendants of u . If G is weakly branching, then G separates the boundary ∂T of the tree, see [1, Proof of Corollary 1.4]. If in addition, the product of rigid stabilizers $\prod_{u \in \mathbb{T}_n} \text{Rist}_G(C_u)$ is a finite index subgroup of G for every n , then G is said to be a *branch group*.

3. A SEQUENCE OF MARKED GROUPS

This section is devoted to the proof of Proposition 1.2. To fix ideas, we start with the Fabrykowski-Gupta group introduced in [14]. It is a group acting on the ternary rooted tree \mathbb{T} . Encode vertices of \mathbb{T} by finite strings in the alphabet $0, 1, 2$, and the boundary of the tree by infinite strings in $0, 1, 2$. Denote by \mathbb{T}_n the level n vertices of the rooted tree \mathbb{T} and $\text{St}_G(n)$ the level n stabilizer, that is, $\text{St}_G(n) = \{g \in G : u \cdot g = u \text{ for all } u \in \mathbb{T}_n\}$.

The Fabrykowski-Gupta group is generated by two elements: a root permutation a which permutes the three subtrees of the root cyclically and a directed permutation b which fixes the right most ray 2^∞ and is defined recursively by

$$b = (a, id, b).$$

In other words we have for any ray $w \in \{0, 1, 2\}^\infty$,

$$\begin{aligned} 0w \cdot a &= 1w, & 1w \cdot a &= 2w, & 2w \cdot a &= 0w; \\ 0w \cdot b &= 0(w \cdot a), & 1w \cdot b &= 1w, & 2w \cdot b &= 2(w \cdot b). \end{aligned}$$

See Figure 3.1. For more background on groups acting on rooted trees and the notation of wreath recursion see the reference [8]. The group $G = \langle a, b \rangle$ is called the Fabrykowski-Gupta group. It is an example of non-torsion Grigorchuk-Gupta-Sidki (GGS) groups.

The group $G = \langle a, b \rangle$ is known to have the following properties:

- ([7]) G is a just infinite branch group which is regularly branching over its commutator group $[G, G]$.
- ([15, 9]) G is of intermediate growth.
- ([16]) G has the congruence subgroup property: every finite index subgroup of G contains some level stabilizer $\text{St}_G(n)$.

3.1. Permutation wreath extensions. Let G_n be the quotient group $G/\text{St}_G(n)$, which acts faithfully and transitively on \mathbb{T}_n . We denote by $\bar{a}, \bar{b} \in G_n$ the images of the generators a, b under the quotient map $G \rightarrow G/\text{St}_G(n)$. Consider the level n Schreier graph \mathcal{S}_n with vertex set \mathbb{T}_n and edge set $E = \{(x, x \cdot \bar{a}), (x, x \cdot \bar{b}) : x \in \mathbb{T}_n\}$. It is a finite graph on 3^n vertices. Consider




FIGURE 3.1. The action of the Fabrykowski-Gupta group on the first four levels of the rooted ternary tree. Self loops are not depicted. Arrows show the action on the roots of subtrees, with corresponding arrows in the rest of the subtree not drawn explicitly. The restriction of \mathcal{S}_3 to its red (and likewise gray and blue) vertices forms a copy of \mathcal{S}_2 . The Schreier graph \mathcal{S}_3 is the disjoint union of these three graphs, with the addition of the three edges labeled by b .

the permutation wreath product of the free product $\mathbf{A} = (\mathbb{Z}/3\mathbb{Z}) * (\mathbb{Z}/3\mathbb{Z})$ and G_n over the set \mathbb{T}_n , that is,

$$\mathbf{A} \wr_{\mathbb{T}_n} G_n = (\oplus_{\mathbb{T}_n} \mathbf{A}) \rtimes G_n,$$

where G_n acts on $\oplus_{\mathbb{T}_n} \mathbf{A}$ by permuting the coordinates. We write elements of $\mathbf{A} \wr_{\mathbb{T}_n} G_n$ as pairs (φ, g) , where $\varphi \in \oplus_{\mathbb{T}_n} \mathbf{A}$ is regarded as a function $\mathbb{T}_n \rightarrow \mathbf{A}$ and $g \in G_n$.

Denote by s and t the two standard generators of \mathbf{A} , $\mathbf{A} = \langle s, t \mid s^3 = t^3 = 1 \rangle$. Consider the subgroup W_n of $\mathbf{A} \wr_{\mathbb{T}_n} G_n$ generated by

$$(3.1) \quad a_n = (id, \bar{a}), \quad b_n = \left(\delta_{2^{n-1}0}^s + \delta_{2^{n-1}1}^{id} + \delta_{2^n}^t, \bar{b} \right),$$

where in the direct sum $\oplus_{\mathbb{T}_n} \mathbf{A}$, δ_x^γ denotes the function that is γ at x and identity elsewhere. We use additive notation $\delta_x^{\gamma_1} + \delta_y^{\gamma_2}$, $x \neq y$, for the function that is γ_1 at x , γ_2 at y , and identity elsewhere.

The choice of (a_n, b_n) guarantees:

Lemma 3.1. *The sequence $(W_n, (a_n, b_n))$ converges to $(G, (a, b))$ in the Cayley-Grigorchuk topology as $n \rightarrow \infty$. Indeed, $(W_{n+1}, (a_{n+1}, b_{n+1}))$ is a marked quotient of $(W_n, (a_n, b_n))$, and the ball of radius 2^{n-2} around id in the Cayley graph of $(W_n, (a_n, b_n))$ coincide with the ball of radius 2^{n-2} around id in $(G, (a, b))$.*

Proof. The Fabrykowski-Gupta group belongs to the class of bounded automaton groups. Schreier graphs of bounded automaton groups are studied

systematically in Bondarenko's dissertation [11]. In particular, on the finite Schreier graph \mathcal{S}_n , we have that the graph distance between the vertices $2^{n-1}0, 2^n$ satisfy $d_{\mathcal{S}_n}(2^{n-1}0, 2^n) = 2^n - 1$. For more details see [11, Chapter VI].

Note that G embeds as a subgroup of $G \wr_{\mathbb{T}_n} G_n$, where the embedding is given by the wreath recursion

$$a \mapsto (id, \bar{a}), \quad b \mapsto \left(\delta_{2^{n-1}0}^a + \delta_{2^{n-1}1}^{id} + \delta_{2^n}^b, \bar{b} \right).$$

Now consider a word $w = w_1 \dots w_\ell$, where $w_j \in \{a^{\pm 1}, b^{\pm 1}\}$ and evaluate this word in $G \wr_{\mathbb{T}_n} G_n$ by the embedding above. Denote the image in $G \wr_{\mathbb{T}_n} G_n$ by (ϕ_w, \bar{w}) . For the configuration $\phi_w \in \oplus_{\mathbb{T}_n} G$, we have that

$$\phi_w(x) = \prod_{i=1}^n \phi_{w_i}(x \cdot w_1 \dots w_{i-1}).$$

It follows from the triangle inequality that if $\ell \leq 2^{n-2}$, then the trajectory $\{x, x \cdot w_1, \dots, x \cdot w_1 \dots w_{\ell-1}\}$ can visit at most one point in the set $\{2^{n-1}0, 2^n\}$. In particular, $\phi_w(x)$ is an element in either $\langle a \rangle$ or $\langle b \rangle$. Thus if we evaluate the same word w in W_n under $a \mapsto a_n$ and $b \mapsto b_n$, the resulting element $(\tilde{\phi}_w, \bar{w})$ can be identified with (ϕ_w, \bar{w}) in $G \wr_{\mathbb{T}_n} G_n$. Namely, ϕ_w is obtained from $\tilde{\phi}_w$ by replacing s with a and t with b and vice versa.

The quotient map from $(W_n, (a_n, b_n))$ to $(W_{n+1}, (a_{n+1}, b_{n+1}))$ is given as follows. Note that $\mathbf{A} \wr_{\mathbb{T}_{n+1}} G_{n+1} = (\mathbf{A} \wr_{\{0,1,2\}} \langle a \rangle) \wr_{\mathbb{T}_n} G_n$. Let $\tau : \mathbf{A} \rightarrow \mathbf{A} \wr_{\{0,1,2\}} \langle a \rangle$ be the group homomorphism determined by $\tau(s) = (id, a)$ and $\tau(t) = (\delta_0^s + \delta_1^{id} + \delta_2^t, id)$. The homomorphism τ extends to $\oplus_{\mathbb{T}_n} \mathbf{A} \rightarrow \oplus_{\mathbb{T}_n} (\mathbf{A} \wr_{\{0,1,2\}} \langle a \rangle)$ coordinate-wise, that is $\tau(\phi)(x) = \tau(\phi(x))$, $x \in \mathbb{T}_n$. It follows that the wreath recursion formula in G that the map

$$\begin{aligned} W_n &\rightarrow W_{n+1} \\ (\phi, g) &\mapsto (\tau(\phi), g) \end{aligned}$$

is a marked group epimorphism which sends a_n to a_{n+1} and b_n to b_{n+1} . \square

Next we verify that W_n is virtually a direct product of free groups:

Lemma 3.2. *The group W_n contains $\oplus_{\mathbb{T}_n} [\mathbf{A}, \mathbf{A}]$ as a finite index normal subgroup.*

Proof. We proceed by induction on n .

As in the proof of Lemma 3.1, let $\tau : \mathbf{A} \rightarrow \mathbf{A} \wr_{\{0,1,2\}} \langle a \rangle$ be the group homomorphism determined by $\tau(s) = (id, a)$ and $\tau(t) = (\delta_0^s + \delta_1^{id} + \delta_2^t, id)$, where a is the 3-cycle $(0, 1, 2)$. When $n = 1$, by definition W_1 is generated by $a_1 = \tau(s)$ and $b_1 = \tau(t)$. Since $a_1^{-1} b_1 a_1 = (\delta_0^{id} + \delta_1^t + \delta_2^s, id)$, it follows that the projection of $W_1 \cap \oplus_{\mathbb{T}_1} \mathbf{A}$ to the component over vertex 2 is \mathbf{A} . Direct calculation shows that $[b_1 a_1^{-1} b_1 a_1, a_1 b_1 a_1^{-1} b_1] = \left(\delta_2^{s t s^{-1} t^{-1}}, id \right)$. It follows

that $[W_1, W_1] \cap \oplus_{\mathbb{T}_1} \mathbf{A}$ contains $\left\{ (\delta_2^\gamma, id) : \gamma \in \langle sts^{-1}t^{-1} \rangle^{\mathbf{A}} \right\}$, while the normal closure $\langle sts^{-1}t^{-1} \rangle^{\mathbf{A}}$ is exactly the commutator subgroup $[\mathbf{A}, \mathbf{A}]$. Since $\tau(s)$ acts as a 3-cycle permuting $\mathbb{T}_1 = \{0, 1, 2\}$, it follows that $[W_1, W_1] \cap \oplus_{\mathbb{T}_1} \mathbf{A} > \oplus_{\mathbb{T}_1} [\mathbf{A}, \mathbf{A}]$. The quotient group $W_1 / \oplus_{\mathbb{T}_1} [\mathbf{A}, \mathbf{A}]$ is a subgroup of $(\mathbf{A} / [\mathbf{A}, \mathbf{A}]) \wr_{\mathbb{T}_1} \langle a \rangle$, which is finite.

We have shown that $\tau([\mathbf{A}, \mathbf{A}])$ contains $\oplus_{\mathbb{T}_1} [\mathbf{A}, \mathbf{A}]$ as finite index normal subgroup, which reflects the property that G is regularly branching over its commutator subgroup.

Suppose the statement is true for n that W_n contains $\oplus_{\mathbb{T}_n} [\mathbf{A}, \mathbf{A}]$ as a finite index normal subgroup. To prove the claim for $n+1$, it suffices to show that $(\delta_{2^{n+1}}^\gamma, id) \in W_{n+1}$ for any $\gamma \in [\mathbf{A}, \mathbf{A}]$. Recall the quotient map $\pi : W_n \rightarrow W_{n+1}$ explained in the proof of Lemma 3.1, where $\mathbf{A} \wr_{\mathbb{T}_{n+1}} G_{n+1}$ is identified with $(\mathbf{A} \wr_{\{0,1,2\}} \langle a \rangle) \wr_{\mathbb{T}_n} G_n$. By the induction hypothesis, $(\delta_{2^n}^\sigma, id) \in W_n$ for any $\sigma \in [\mathbf{A}, \mathbf{A}]$. Under the quotient map π , we have

$$\pi((\delta_{2^n}^\sigma, id)) = (\delta_{2^n}^{\tau(\sigma)}, id).$$

With the map τ we are back in the situation of the induction base, where we have shown that $\tau([\mathbf{A}, \mathbf{A}])$ contains $\oplus_{\mathbb{T}_1} [\mathbf{A}, \mathbf{A}]$. In particular, for any $\gamma \in [\mathbf{A}, \mathbf{A}]$, there is an element $\sigma \in [\mathbf{A}, \mathbf{A}]$ such that $\tau(\sigma) = (\delta_2^\gamma, id)$. It follows that $\pi((\delta_{2^n}^\sigma, id)) = (\delta_{2^n}^{\tau(\sigma)}, id) = (\delta_{2^{n+1}}^\gamma, id)$, in particular it is an element of W_{n+1} . □

3.2. Choices of marked subgroups. We are given a fixed rank $d \in \mathbb{N}$, $d \geq 2$. Since the Fabrykowski-Gupta group $G = \langle a, b \rangle$ is a branch group acting faithfully on the ternary tree \mathbb{T} , by Abért's criterion [1, Theorem 1.1] and its proof, we have that given any $n \in \mathbb{N}$, and any vertex $v \in \mathbb{T}$, there exist elements $\gamma_1^{(n)}, \dots, \gamma_d^{(n)} \in \text{Rist}_G(v)$ such that $w \left(a\gamma_1^{(n)}, b\gamma_2^{(n)}, \gamma_3^{(n)}, \dots, \gamma_d^{(n)} \right) \neq id$ for all reduced word w of length $1 \leq |w| \leq n$.

In what follows we fix the choice of v to be the child of the root indexed by 1. For each $n \in \mathbb{N}$, fix a choice of $\gamma_1^{(n)}, \dots, \gamma_d^{(n)} \in \text{Rist}_G(1)$ such that the tuple $(a\gamma_1^{(n)}, b\gamma_2^{(n)}, \gamma_3^{(n)}, \dots, \gamma_d^{(n)})$ do not satisfy any reduced word w of length $|w| \in [1, n]$.

Lemma 3.3. *Denote by H_n the subgroup of G generated by the first two elements of the tuple chosen above,*

$$H_n = \langle a\gamma_1^{(n)}, b\gamma_2^{(n)} \rangle.$$

Then H_n acts level transitively on the rooted ternary tree \mathbb{T} .

Proof. The statement is equivalent to that the Schreier graph on level k vertices \mathbb{T}_k with respect to $(a\gamma_1^{(n)}, b\gamma_2^{(n)})$ is connected.

With respect to the original generators a, b , the Schreier graph \mathcal{S}_k with vertex set \mathbb{T}_k under the action of G can be drawn recursively as follows, see [11, Chapter V]. The level 1 graph \mathcal{S}_1 on $\{0, 1, 2\}$ has (directed) edges $(0, 1)$, $(1, 2)$, $(2, 0)$ labeled by a , and self loops at each vertex labeled by b . To draw \mathcal{S}_{k+1} , take three copies of \mathcal{S}_k , append letter 0, 1, or 2 to the strings indexing the vertices of \mathcal{S}_k respectively in each copy. Then connect the three copies by edges $(2^{k-1}00, 2^{k-1}01)$, $(2^{k-1}01, 2^{k-1}02)$ and $(2^{k-1}02, 2^{k-1}00)$ labeled by b . In Figure 3.1 this is depicted for $k = 2$. There, \mathcal{S}_3 is seen to be the union of three copies of \mathcal{S}_2 , which are shown with red, gray and blue vertices, respectively. The three additional edges connecting them and labeled by b are those in the bottom right of the figure.

It follows that in \mathcal{S}_k , if we remove all vertices of the form $1u$, $u \in \{0, 1, 2\}^{k-1}$ and edges connecting to such vertices, the remaining graph is connected. Denote the remaining graph by \mathcal{S}'_k . Since $\gamma_1^{(n)}, \gamma_2^{(n)}$ are chosen to be in the rigid stabilizer of vertex 1, in the subgraph \mathcal{S}'_k , we may replace label a by $a\gamma_1^{(n)}$ and label b by $b\gamma_2^{(n)}$. The element $a\gamma_1^{(n)}$ moves $1u$, $u \in \{0, 1, 2\}^{k-1}$ into the vertex set of \mathcal{S}'_k , namely, a string starting with 2. It follows that the graph on \mathbb{T}_k with respect to $(a\gamma_1^{(n)}, b\gamma_2^{(n)})$ is connected. \square

Denote by ℓ_n the maximal word length of elements in the tuple chosen, with respect to the original generating set (a, b) , that is,

$$(3.2) \quad \ell_n = \max \left\{ \left| a\gamma_1^{(n)} \right|_{\{a,b\}}, \left| b\gamma_2^{(n)} \right|_{\{a,b\}}, \dots, \left| \gamma_d^{(n)} \right|_{\{a,b\}} \right\}.$$

Recall the subgroup W_k of $\mathbf{A} \wr_{\mathbb{T}_k} G_k$, generated by (a_k, b_k) , as defined in (3.1). By Lemma 3.1, the ball of radius 2^{k-2} around the identity element in the Cayley graph of $(W_k, (a_k, b_k))$ coincide with the ball of radius 2^{k-2} around the identity in $(G, (a, b))$. In particular, for $k \geq 2 + \log_2(n\ell_n)$, elements in the tuple $(a\gamma_1^{(n)}, b\gamma_2^{(n)}, \gamma_3^{(n)}, \dots, \gamma_d^{(n)})$ have images in $(W_k, (a_k, b_k))$ under the identification of balls of radius $2\ell_n$ around the identities. Record the image tuple of elements in W_k as $(h_1^{(n,k)}, \dots, h_d^{(n,k)})$. Finally, denote by $\Gamma_{n,k}$ the subgroup of W_k generated by the tuple $S_{n,k} = (h_1^{(n,k)}, \dots, h_d^{(n,k)})$. Note that by our choices, for $k \geq 2 + \log_2(n\ell_n)$, the ball of radius n around the identity in the Cayley graph of $(\Gamma_{n,k}, S_{n,k})$ is the same as the ball of radius n in the Cayley graph of the free group $(\mathbf{F}_d, \mathbf{S})$.

3.3. Random walks on $\Gamma_{n,k}$. Let μ be a non-degenerate probability measure on the free group \mathbf{F}_d . Next we consider the μ -random walk on the group $\Gamma_{n,k}$ defined in the previous subsection, with $n \geq 1$, $k \gg \log_2(n\ell_n)$. Our goal is to show that the action of $\Gamma_{n,k}$ on the Poisson boundary of $(\Gamma_{n,k}, \mu)$ is essentially free; and given any $\epsilon > 0$, for all k sufficiently large, the asymptotic entropy of the μ -random walk on $\Gamma_{n,k}$ is smaller than ϵ .

By [13, Theorem 5.1], for a non-degenerate probability measure μ on a countable group Γ , a sufficient condition for the action of Γ on the Poisson boundary of (Γ, μ) to be essentially free is that

- (1): Γ has only countably many amenable subgroups,
- (2): Γ does not contain any non-trivial normal amenable subgroup, in other words, the amenable radical of Γ is trivial.

We verify that these two properties are satisfied by $\Gamma_{n,k}$ in the following lemma.

Lemma 3.4. *For each n and $k \geq 2 + \log_2(n\ell_n)$, the group $\Gamma_{n,k}$ is non-amenable, has no non-trivial normal amenable subgroup, and has only countably many amenable subgroups.*

Proof. We first introduce some notations. Given a subgroup H of $\mathbf{A} \wr_{\mathbb{T}_k} G_k = (\oplus_{\mathbb{T}_k} \mathbf{A}) \rtimes G_k$, for $g \in G_k$, let

$$S_H(g) := \{\phi \in \oplus_{\mathbb{T}_k} \mathbf{A} : (\phi, g) \in H\}.$$

Then $S_H(id_{G_k})$ is a subgroup of $\oplus_{\mathbb{T}_k} \mathbf{A}$. For $g \in G_k$, $S_H(g)$ is either empty, or a right coset of $S_H(id_{G_k})$ in $\oplus_{\mathbb{T}_k} \mathbf{A}$. Denote by π_k the natural projection $\mathbf{A} \wr_{\mathbb{T}_k} G_k \rightarrow G_k$.

As shown in the proof of Lemma 3.3, in the Schreier graph on vertex set \mathbb{T}_k with respect to the generating tuple of H_n , there is a path connecting the vertex 2^k to the vertex $2^{k-1}0$, which does not visit any vertex starting with letter 1. It then follows by the definition of the generating tuple $S_{n,k} = (h_1^{(n,k)}, \dots, h_d^{(n,k)})$ of $\Gamma_{n,k}$ that the set of values at the coordinate index by 2^k is all \mathbf{A} , that is,

$$\{\phi(2^k) : \phi \in \cup_{g \in G_k} S_{\Gamma_{n,k}}(g)\} = \mathbf{A}$$

Note that this implies that $S_{\Gamma_{n,k}}(id_{G_k})$ is non-amenable. Indeed, otherwise the free product \mathbf{A} can be written as a union of finitely many right cosets of an amenable subgroup, which contradicts the fact that \mathbf{A} is non-amenable. For each $x \in \mathbb{T}_k$, write θ_x for the projection of $\oplus_{\mathbb{T}_k} \mathbf{A}$ to the x -coordinate, that is, $\theta_x(\phi) = \phi(x)$. Then the reasoning above shows that the projection of $\Gamma_{n,k} \cap (\oplus_{\mathbb{T}_k} [\mathbf{A}, \mathbf{A}])$ under θ_{2^k} is non-amenable. Recall that $[\mathbf{A}, \mathbf{A}]$ is a free group, By Lemma 3.3, the action of $\pi_k(\Gamma_{n,k})$ is transitive on \mathbb{T}_k . It follows then $\theta_x(\Gamma_{n,k} \cap (\oplus_{\mathbb{T}_k} [\mathbf{A}, \mathbf{A}]))$ is a free group of rank at least 2 for every vertex in \mathbb{T}_k .

Let N be a normal subgroup of $\Gamma_{n,k}$, $N \neq \{id\}$. We need to show that N is non-amenable. Note that for each $x \in \mathbb{T}_k$, $\theta_x(N \cap \oplus_{\mathbb{T}_k} [\mathbf{A}, \mathbf{A}])$ a normal subgroup of $\theta_x(\Gamma_{n,k} \cap (\oplus_{\mathbb{T}_k} [\mathbf{A}, \mathbf{A}]))$, while the latter is a free group of rank at least 2. Thus if on the contrary N is amenable, then $N \cap \oplus_{\mathbb{T}_k} [\mathbf{A}, \mathbf{A}] = \{id\}$. Since $\mathbf{A}/[\mathbf{A}, \mathbf{A}]$ is finite, such trivial intersection implies that $S_N(id_{G_k})$ is finite.

We now argue that $S_N(id_{G_k})$ being a finite group contradicts with the condition that N is a non-trivial normal subgroup of $\Gamma_{n,k}$. Note that since

G_k is finite, $S_N(id_{G_k})$ being finite implies that N is finite. On the other hand, since for each $x \in \mathbb{T}_k$, $\theta_x(\Gamma_{n,k} \cap (\oplus_{\mathbb{T}_k}[\mathbf{A}, \mathbf{A}]))$ is a free group of rank at least 2, it follows that for any element $h \in \Gamma_{n,k}$, $h \neq id$, its conjugacy class is infinite. Therefore $S_N(id_{G_k})$ being a finite group implies that $N = \{id\}$. We conclude that a nontrivial normal subgroup of $\Gamma_{n,k}$ is non-amenable.

Since $[\mathbf{A}, \mathbf{A}]$ is a free group, the only amenable subgroups are the trivial group and the cyclic groups. It follows that the direct sum $\oplus_{\mathbb{T}_n}[\mathbf{A}, \mathbf{A}]$ has only countably many amenable subgroups. The property of having only countably many amenable subgroups is clearly preserved under taking finite extensions and taking subgroups. Thus by Lemma 3.2, $\Gamma_{n,k}$ has only countably many amenable subgroups. \square

To bound the asymptotic entropy from above, we simply use the well-known ‘‘fundamental inequality’’, see e.g. [10]. More precisely, let μ be a probability measure on \mathbf{F}_d with finite first moment, $\pi : \mathbf{F}_d \rightarrow \Gamma$ an epimorphism. Let $S = \pi(\mathbf{S})$ be the induced marking on Γ and $\bar{\mu} = \pi \circ \mu$ be the pushforward of μ . The fundamental inequality implies that

$$h(\Gamma, \bar{\mu}) \leq v_{\Gamma, S} \cdot \ell_{\Gamma, \bar{\mu}},$$

where $v_{\Gamma, S}$ and $\ell_{\Gamma, \bar{\mu}}$ are asymptotic volume growth rate and asymptotic speed with respect to generating set S :

$$v_{\Gamma, S} = \lim_{r \rightarrow \infty} \frac{1}{r} \log V_{\Gamma, S}(r) \text{ and } \ell_{\Gamma, \bar{\mu}} = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{g \in \Gamma} |g|_S \bar{\mu}^{(n)}(g).$$

By sub-additivity, we have $\ell_{\Gamma, \bar{\mu}} \leq \sum_{g \in \Gamma} |g|_S \bar{\mu}(g) \leq \sum_{g \in \mathbf{F}_d} |g|_S \mu(g)$. Thus the asymptotic entropy can be bounded by

$$(3.3) \quad h(\Gamma, \bar{\mu}) \leq v_{\Gamma, S} \sum_{g \in \mathbf{F}_d} |g|_S \mu(g).$$

The estimate (3.3) is the only place where the moment condition on μ is needed.

By Lemma 3.1, the ball of radius 2^{k-2} around id in the Cayley graph of $(W_k, (a_k, b_k))$ coincide with the ball of radius 2^{k-2} around id in $(G, (a, b))$. It follows by sub-multiplicity of the volume growth function that the asymptotic volume rate satisfy

$$v_{W_k, (a_k, b_k)} \leq \frac{1}{2^{k-2}} \log V_{G, (a, b)}(2^{k-2}).$$

Recall the maximal length ℓ_n defined in (3.2). By comparing lengths of generators, we have that for the subgroup $\Gamma_{n,k}$ of W_k satisfies

$$v_{\Gamma_{n,k}, S_{n,k}} \leq \ell_n v_{W_k, (a_k, b_k)}.$$

3.4. Proof of Proposition 1.2. We are now ready to prove Proposition 1.2 stated in the Introduction.

Completion of proof of Proposition 1.2. Let μ be a non-degenerate probability measure on \mathbf{F}_d given, where $d \geq 2$ and μ is of finite first moment. For each $n \in \mathbb{N}$, choose $k_n \gg \log_2(n\ell_n)$ such that Γ_{n,k_n} is defined as in subsection 3.2 and moreover

$$\frac{\ell_n}{2^{k_n-2}} \log V_{G,(a,b)}(2^{k_n-2}) \leq \frac{1}{n}.$$

This is possible because the Fabrykowski-Gupta group G has sub-exponential volume growth, that is,

$$v_{G,(a,b)} = \lim_{r \rightarrow \infty} \frac{1}{r} \log V_{G,(a,b)}(r) = 0,$$

Now we verify that the sequence $(\Gamma_{n,k_n}, S_{n,k_n})$ satisfy the properties stated.

- (i): By construction, the generating tuple S_{n,k_n} do not satisfy any reduced word w of length $|w| \in [1, n]$.
- (ii): The fundamental inequality (3.3) implies that with respect to the marking $(\mathbf{F}_d, \mathbf{S}) \rightarrow (\Gamma_{n,k_n}, S_{n,k_n})$,

$$h(\Gamma_{n,k_n}, \mu) \leq v_{\Gamma_{n,k_n}, S_{n,k_n}} \sum_{g \in \mathbf{F}_d} |g|_{\mathbf{S}} \mu(g) \leq \frac{1}{n} \sum_{g \in \mathbf{F}_d} |g|_{\mathbf{S}} \mu(g).$$

Thus the sequence of asymptotic entropies converge to 0 as $n \rightarrow \infty$.

- (iii): This property is shown in Lemma 3.4.

The proof of Proposition 1.2 is complete. □

Remark 3.5. For $d \geq 3$, in the proof of Proposition 1.2 one can use the first Grigorchuk group $G_{012} = \langle a, b, c \rangle$ introduced in [22, 23] instead of the Fabrykowski-Gupta group. Recall that G_{012} acts on the rooted binary tree. Then one can consider the permutational wreath extension $\mathbf{B} \wr_{\mathbf{T}_n} G_n$, where $G_n = G/\text{St}_G(n)$ and $\mathbf{B} = (\mathbb{Z}/2\mathbb{Z}) * (\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}) = (\langle \mathbf{s} \rangle) * (\langle \mathbf{t} \rangle \times \langle \mathbf{u} \rangle)$. Similar to the sequence of extensions Γ_n , set

$$H_n = \langle a_n, b_n, c_n \rangle < \mathbf{B} \wr_{\mathbf{T}_{3n}} G_{3n}$$

where the generators are defined as

$$a_n = (id, \bar{a}), \quad b_n = (\delta_{1^{3n}}^{\mathbf{t}} + \delta_{1^{3n-1}0}^{\mathbf{s}}, \bar{b}), \quad c_n = (\delta_{1^{3n}}^{\mathbf{u}} + \delta_{1^{3n-1}0}^{\mathbf{s}}, \bar{c}).$$

Similar proof as in this section with Γ_n replaced by H_n shows that for $d \geq 3$, the statement Proposition 1.2 is true under the weaker assumption that μ has finite α_0 -moment and finite entropy, where α_0 is the exponent in the growth upper bound $v_{G_{012}}(r) \lesssim e^{r^{\alpha_0}}$ from [5, 35], $\alpha_0 \approx 0.7674$.

We choose to take extensions of the Fabrykowski-Gupta group G here because the resulting groups are 2-generated, which allows to cover the case $d = 2$. It is remarked in [17] that all maximal subgroups of G are of finite index, which would imply that there is a sequence of marking S_k on G such that (G, S_k) converges to the free group $(\mathbf{F}_d, \mathbf{S})$ when $k \rightarrow \infty$ by [6]. Since

we could not find a written proof of this statement, in this section we produce tuples of elements of G , which a priori do not necessarily generate G , where only Abért's criterion [1] is invoked.

4. STATIONARY JOININGS AND PROOF OF THE MAIN RESULTS

Let (X, ν) and (Y, η) be two μ -stationary G -spaces. Following [21], we say a probability measure λ on $X \times Y$ is a stationary joining of ν and η if it is μ -stationary and its marginals are ν and η respectively.

In this section we focus on the situation where both stationary systems are (G, μ) -boundaries. We use notations introduced in Section 2.1. Denote by (B, ν_B) a compact model of the Poisson boundary of (G, μ) . Let (X, ν) and (Y, η) be compact models of two (G, μ) -boundaries and denote by β_X and β_Y the corresponding maps from the Poisson boundary (B, ν_B) to (X, ν) and (Y, η) . Consider the map

$$\begin{aligned} \beta_X \times \beta_Y : B &\rightarrow X \times Y \\ b &\mapsto (\beta_X(b), \beta_Y(b)), \end{aligned}$$

and denote by Z the range $(\beta_X \times \beta_Y)(B)$ and $\nu \vee \eta$ the pushforward of the harmonic measure ν_B under $\beta_X \times \beta_Y$. Then it's clear by definition that $(Z, \nu \vee \eta)$ is a G -factor of the Poisson boundary (B, ν_B) , in other words, it is a (G, μ) -boundary. The G -space $(Z, \nu \vee \eta)$ is the unique stationary joining of the μ -boundaries (X, ν) and (Y, η) , see [21, Proposition 3.1].

On the level of groups, given two d -marked groups (G_1, S_1) and (G_2, S_2) , one can take their diagonal product, denoted by $(G_1 \otimes G_2, S)$, as the subgroup of $G_1 \times G_2$ generated by

$$S = \left(\left(s_1^{(1)}, s_1^{(2)} \right), \dots, \left(s_d^{(1)}, s_d^{(2)} \right) \right),$$

where $S_i = \left(s_1^{(i)}, \dots, s_d^{(i)} \right)$, $i = 1, 2$. This operation on two groups corresponds to taking stationary joinings of the Poisson boundaries:

Lemma 4.1. *Let μ be a probability measure on \mathbf{F}_d . The Poisson boundary of $(G_1 \otimes G_2, \mu)$ is the stationary joining of the Poisson boundaries of (G_1, μ) and (G_2, μ) .*

Proof. Denote by (B_i, ν_i) the Poisson boundary of (G_i, μ) , $i = 1, 2$ and regard them as $G_1 \otimes G_2$ -spaces. Denote by $(Z, \nu_1 \vee \nu_2)$ the stationary joining of (B_1, ν_1) and (B_2, ν_2) as above and $\pi_i : Z \rightarrow B_i$ the projections. We need to show $(Z, \nu_1 \vee \nu_2)$ is the maximal $(G_1 \otimes G_2, \mu)$ -boundary.

Let (Y, η) be a $(G_1 \otimes G_2, \mu)$ -boundary. Denote by K_i the subgroup of $G_1 \otimes G_2$ which consists of elements that project to identity in G_i , that is,

$$K_i = \{(g_1, g_2) \in G_1 \times G_2 : (g_1, g_2) \in G_1 \otimes G_2, g_i = id_{G_i}\}.$$

Denote by $Y_i = Y/K_i$ the space of K_i -ergodic components of Y and η_i the pushforward of the measure η under the K_i -factor map $Y \rightarrow Y/K_i$. Since (Y, η) is an ergodic $G_1 \otimes G_2$ -space and $K_1 \cap K_2 = \{id\}$, we have that Y can

be viewed as a subset of $Y_2 \times Y_1$. It's easy to see that by definition of K_i that $G_1 \otimes G_2/K_2 \simeq G_1$. It follows that (Y_2, η_2) is a (G_1, μ) -boundary. Denote by β_{Y_2} the boundary map from (B_1, ν_1) to (Y_2, η_2) . In the same way we have (Y_1, η_1) is a (G_2, μ) -boundary and denote by $\beta_{Y_1} : (B_2, \nu_2) \rightarrow (Y_1, \eta_1)$ the boundary map. By uniqueness of stationary joinings of μ -boundaries, we have that $(Y, \eta) = (Y_2 \times Y_1, \eta_2 \vee \eta_1)$. It follows that (Y, η) is a factor of $(Z, \nu_1 \vee \nu_2)$, where the boundary map is given by $z \mapsto (\beta_{Y_2} \circ \pi_1(z), \beta_{Y_1} \circ \pi_2(z))$. \square

With the sequence of marked groups provided by Proposition 1.2, we are now ready to complete the proofs of Theorem 1.1 and 1.3.

Proof of Theorem 1.1. Denote by (B, ν_B) the Poisson boundary of (\mathbf{F}_d, μ) . Let $((\Gamma_k, S_k))_{k=1}^\infty$ be a sequence marked groups provided by Proposition 1.2. Denote by (Π_k, η_k) the Poisson boundary of (Γ_k, μ) . Since (Γ_k, S_k) can be identified with a projection $\pi_k : \mathbf{F}_d \rightarrow \Gamma_k$, we regard (Π_k, η_k) as a (\mathbf{F}_d, μ) -space, where the \mathbf{F}_d -action factors through π_k .

Since Γ is a proper quotient of \mathbf{F}_d , $N = \ker(\pi : \mathbf{F}_d \rightarrow \Gamma)$ is nontrivial. Fix a choice of element $g \in N$, $g \neq id$. Choose an index $k \in \mathbb{N}$ sufficiently large such that the balls of radius $2|g|_{\mathbf{S}}$ around identities in (Γ_k, S_k) and $(\mathbf{F}_d, \mathbf{S})$ coincide and $h(\Gamma_k, \mu) < \epsilon$. Take $\tilde{\Gamma}$ to be the diagonal product $(\Gamma \otimes \Gamma_k, S)$. Then

$$h(\Gamma \otimes \Gamma_k, \mu) \leq h(\Gamma, \mu) + h(\Gamma_k, \mu) < h(\Gamma, \mu) + \epsilon.$$

Since g acts trivially on the Poisson boundary of (Γ, μ) but acts freely on (Π_k, ν_k) , it follows that (Π_k, ν_k) is not a \mathbf{F}_d -factor of the Poisson boundary of (Γ, μ) . By Lemma 2.2, we conclude that $h(\Gamma \otimes \Gamma_k, \mu) > h(\Gamma, \mu)$. \square

Proof of Theorem 1.3. The proof is similar to Theorem 1.1. Since (X, ν) is assumed to be a (\mathbf{F}_d, μ) -boundary where the action of \mathbf{F}_d is not essentially free, we can choose an element $g \in \mathbf{F}_d$, $g \neq 1$, such that $\nu(\text{Fix}_X(g)) > 0$. Choose an index $k \in \mathbb{N}$ sufficiently large such that the balls of radius $2|g|_{\mathbf{S}}$ around identities in (Γ_k, S_k) and $(\mathbf{F}_d, \mathbf{S})$ coincide and $h(\Gamma_k, \mu) < \epsilon$. Take the stationary joining $(Z_k, \nu \vee \eta_k)$ of (X, ν) and (Π_k, η_k) . By the general inequality, we have

$$h(Z_k, \nu \vee \eta_k) \leq h(X, \nu) + h(\Pi_k, \eta_k) \leq h(X, \nu) + \epsilon.$$

It remains to show that $h(Z_k, \nu \vee \eta_k) > h(X, \nu)$. Suppose on the contrary equality holds, then by Lemma 2.2, the equality would imply $(Z_k, \nu \vee \eta_k) = (X, \nu)$. However the action of Γ_k on (Π_k, η_k) is essentially free, which implies $\nu \vee \eta_k(\text{Fix}_{Z_k}(g)) = 0$, contradicting $\nu(\text{Fix}_X(g)) > 0$. \square

We now show an analogous result on spectral radii stated as Theorem 1.4 in the Introduction. Consider a symmetric non-degenerate probability measure μ on Γ . In [32, 33] Kesten proved the following theorem: let μ be

a symmetric non-degenerate probability measure on Γ and N be a normal subgroup of Γ , then the following are equivalent:

- (i): $\rho(\Gamma, \mu) = \rho(\Gamma/N, \mu)$,
- (ii): N is amenable.

Given a proper quotient Γ of \mathbf{F}_d and $\epsilon > 0$, to prove Theorem 1.4 we take $\tilde{\Gamma}$ to be a diagonal product $\Gamma \otimes H$, for some appropriate choice of H similar to the groups used in Theorem 1.1.

Proof of Theorem 1.4. Let $\epsilon > 0$ be a constant given. Let Γ be a proper quotient of \mathbf{F}_d and fix a choice of $g_0 \in \ker(\mathbf{F}_d \rightarrow \Gamma)$, $g_0 \neq id$. Take $n \geq 2|g_0|_{\mathbf{S}}$. As in the beginning of Subsection 3.2, fix a choice of d -tuple of elements in the Fabrykowski-Gupta group G , $T = (a\gamma_1^{(n)}, b\gamma_2^{(n)}, \gamma_3^{(n)}, \dots, \gamma_d^{(n)})$ that do not satisfy any reduced word of length at most n . Denote the group generated by this tuple by G_n . Take first the diagonal product $\Gamma \otimes G_n$. By the choice of g and marking on G we have that $N_0 = \ker(\Gamma \otimes G_n \rightarrow \Gamma)$ is non-trivial. Note that N_0 can be regarded as a normal subgroup of G_n .

Denote by (W_ℓ) a μ -random walk on \mathbf{F}_d . For a marked group (H, S) , we write π_H for the quotient map $\mathbf{F}_d \rightarrow H$ when the marking is clear from the context.

Take a small constant $\epsilon_1 > 0$, choose ℓ large enough such that

$$\mathbb{P}(\pi_\Gamma(W_\ell) = id_\Gamma) \geq ((1 - \epsilon_1)\rho(\Gamma, \mu))^\ell.$$

For $g \in N_0$, set

$$Q(g) = \frac{\mathbb{P}(\pi_{\Gamma \otimes G}(W_\ell) = g)}{\mathbb{P}(\pi_\Gamma(W_\ell) = id_\Gamma)}.$$

Then Q is a symmetric probability measure on N_0 . Equip N_0 with the induced metric $|\cdot|_T$ from (G_n, T) . Let R be a sufficiently large radius such that $Q(\{\gamma \in N_0 : |\gamma|_T > R\}) \leq \epsilon_1$. Truncate the measure Q at R and let

$$Q_R(g) = \frac{1}{Q(\{\gamma : |\gamma|_T \leq R\})} Q(g) \mathbf{1}_{\{|\gamma|_T \leq R\}}.$$

Since N_0 is a subgroup of the Fabrykowski-Gupta group G , thus amenable, there exists an integer m such that

$$Q_R^{2m}(id_{N_0}) \geq (1 - \epsilon_1)^{2m}.$$

With ℓ, m, R chosen as above, for a sufficiently large index k , to be specified shortly, and take the marked group $(\Gamma_{n,k}, S_{n,k})$ defined in Subsection 3.2. Consider the diagonal product $\Gamma \otimes \Gamma_{n,k}$. By Lemma 3.1, the ball of radius 2^{k-2} around id in the Cayley graph of $(W_k, (a_k, b_k))$ coincide with the ball of radius 2^{k-2} around id in $(G, (a, b))$. Choose k sufficiently large such that $k > 2mR$.

Now we follow the original argument in Kesten's theorem (ii) \Rightarrow (i) above to show $\rho(\Gamma \otimes \Gamma_{n,k}, \mu) > \rho(\Gamma, \mu) - \epsilon$. Write $W_{(k-1)\ell}^{k\ell} = W_{(k-1)\ell}^{-1} W_{k\ell}$.

$$\begin{aligned} & \mathbb{P}(\pi_{\Gamma \otimes \Gamma_{n,k}}(W_{2\ell m}) = id_{\Gamma \otimes \Gamma_{n,k}}) \\ & \geq \mathbb{P}\left(\bigcap_{k=1}^{2m} \left\{ \pi_{\Gamma}\left(W_{(k-1)\ell}^{k\ell}\right) = id_{\Gamma}, \left| \pi_{\Gamma_{n,k}}\left(W_{(k-1)\ell}^{k\ell}\right) \right|_{S_{n,k}} \leq R \right\} \cap \left\{ \pi_{\Gamma_{\ell}}(W_{2\ell m}) = id_{\Gamma_{n,k}} \right\}\right) \\ & \geq ((1 - \epsilon_1)\rho(\Gamma, \mu))^{2\ell m} (1 - \epsilon_1)^{2m} Q_R^{2m}(id_{N_0}) \geq (1 - \epsilon_1)^{2mn+4m} \rho^{2mn}. \end{aligned}$$

Choose $\epsilon_1 < \epsilon/3$, we have that $\rho(\Gamma \otimes \Gamma_{n,k}, \mu) > (1 - \epsilon)\rho(\Gamma, \mu)$.

Finally, by Lemma 3.4, $\Gamma_{n,k}$ has no nontrivial amenable normal subgroups. Since by the choice of markings $\ker(\Gamma \otimes G_{n,k} \rightarrow \Gamma)$ is nontrivial, the kernel is non-amenable. By Kesten's theorem (i) \Rightarrow (ii), we conclude that $\rho(\tilde{\Gamma}, \mu) < \rho(\Gamma, \mu)$. □

REFERENCES

- [1] M. Abért. Group laws and free subgroups in topological groups. *Bull. London Math. Soc.*, 37(4):525–534, 2005.
- [2] A. Akhmedov. On the girth of finitely generated groups. *J. Algebra*, 268(1):198–208, 2003.
- [3] A. Avez. Théorème de Choquet-Deny pour les groupes à croissance non exponentielle. *CR Acad. Sci. Paris Sér. A*, 279:25–28, 1974.
- [4] U. Bader and Y. Shalom. Factor and normal subgroup theorems for lattices in products of groups. *Invent. Math.*, 163(2):415–454, 2006.
- [5] L. Bartholdi. The growth of Grigorchuk's torsion group. *Internat. Math. Res. Notices*, (20):1049–1054, 1998.
- [6] L. Bartholdi and A. Erschler. Ordering the space of finitely generated groups. *Ann. Inst. Fourier (Grenoble)*, 65(5):2091–2144, 2015.
- [7] L. Bartholdi and R. I. Grigorchuk. On parabolic subgroups and Hecke algebras of some fractal groups. *Serdica Math. J.*, 28(1):47–90, 2002.
- [8] L. Bartholdi, R. I. Grigorchuk, and Z. Šunić. Branch groups. In *Handbook of algebra, Vol. 3*, volume 3 of *Handb. Algebr.*, pages 989–1112. Elsevier/North-Holland, Amsterdam, 2003.
- [9] L. Bartholdi and F. Pochon. On growth and torsion of groups. *Groups Geom. Dyn.*, 3(4):525–539, 2009.
- [10] S. Blachère, P. Haïssinsky, and P. Mathieu. Asymptotic entropy and Green speed for random walks on countable groups. *Ann. Probab.*, 36(3):1134–1152, 2008.
- [11] I. Bondarenko. *Groups generated by bounded automata and their Schreier graphs*. Texas A&M University, 2007.
- [12] L. Bowen. Random walks on random coset spaces with applications to Furstenberg entropy. *Invent. Math.*, 196(2):485–510, 2014.
- [13] L. Bowen, Y. Hartman, and O. Tamuz. Generic stationary measures and actions. *Trans. Amer. Math. Soc.*, 369(7):4889–4929, 2017.
- [14] J. Fabrykowski and N. Gupta. On groups with sub-exponential growth functions. *J. Indian Math. Soc. (N.S.)*, 49(3-4):249–256 (1987), 1985.
- [15] J. Fabrykowski and N. Gupta. On groups with sub-exponential growth functions. II. *J. Indian Math. Soc. (N.S.)*, 56(1-4):217–228, 1991.
- [16] G. A. Fernández-Alcober, A. Garrido, and J. Uria-Albizuri. On the congruence subgroup property for GGS-groups. *Proc. Amer. Math. Soc.*, 145(8):3311–3322, 2017.

- [17] D. Francoeur and A. Garrido. Maximal subgroups of groups of intermediate growth. *Adv. Math.*, 340:1067–1107, 2018.
- [18] H. Furstenberg. Noncommuting random products. *Transactions of the American Mathematical Society*, 108(3):377–428, 1963.
- [19] H. Furstenberg. Random walks and discrete subgroups of Lie groups. *Advances in Probability and Related Topics*, 1:1–63, 1971.
- [20] H. Furstenberg. Boundary theory and stochastic processes on homogeneous spaces. *Harmonic analysis on homogeneous spaces*, 26:193–229, 1973.
- [21] H. Furstenberg and E. Glasner. Stationary dynamical systems. In *Dynamical numbers—interplay between dynamical systems and number theory*, volume 532 of *Contemp. Math.*, pages 1–28. Amer. Math. Soc., Providence, RI, 2010.
- [22] R. I. Grigorchuk. On Burnside’s problem on periodic groups. *Funktsional. Anal. i Prilozhen.*, 14(1):53–54, 1980.
- [23] R. I. Grigorchuk. Degrees of growth of finitely generated groups and the theory of invariant means. *Izv. Akad. Nauk SSSR Ser. Mat.*, 48(5):939–985, 1984.
- [24] R. I. Grigorchuk. Just infinite branch groups. In *New horizons in pro-p groups*, volume 184 of *Progr. Math.*, pages 121–179. Birkhäuser Boston, Boston, MA, 2000.
- [25] Y. Hartman and O. Tamuz. Furstenberg entropy realizations for virtually free groups and lamplighter groups. *Journal d’Analyse Mathématique*, 126(1):227–257, 2015.
- [26] Y. Hartman and A. Yadin. Furstenberg entropy of intersectional invariant random subgroups. *Compositio Mathematica*, 154(10):2239–2265, 2018.
- [27] V. A. Kaimanovich. Double ergodicity of the poisson boundary and applications to bounded cohomology. *Geometric & Functional Analysis*, 13(4):852–861, 2003.
- [28] V. A. Kaimanovich. Amenability and the Liouville property. *Israel Journal of Mathematics*, 149(1):45–85, 2005.
- [29] V. A. Kaimanovich and A. M. Vershik. Random walks on discrete groups: boundary and entropy. *Ann. Probab.*, 11(3):457–490, 1983.
- [30] M. Kassabov and I. Pak. Groups of oscillating intermediate growth. *Ann. of Math. (2)*, 177(3):1113–1145, 2013.
- [31] M. Kassabov and I. Pak. Unpublished manuscript. *Results announced in Kassabov’s lecture*, <https://www.msri.org/workshops/770/schedules/21638>.
- [32] H. Kesten. Full banach mean values on countable groups. *Mathematica Scandinavica*, pages 146–156, 1959.
- [33] H. Kesten. Symmetric random walks on groups. *Transactions of the American Mathematical Society*, 92(2):336–354, 1959.
- [34] G. W. Mackey. Point realizations of transformation groups. *Illinois Journal of Mathematics*, 6(2):327–335, 1962.
- [35] R. Muchnik and I. Pak. On growth of Grigorchuk groups. *Internat. J. Algebra Comput.*, 11(1):1–17, 2001.
- [36] A. Nevo. The spectral theory of amenable actions and invariants of discrete groups. *Geometriae Dedicata*, 100(1):187–218, 2003.
- [37] A. Nevo and R.J. Zimmer. Rigidity of Furstenberg entropy for semisimple Lie group actions. *Annales Scientifiques de l’École Normale Supérieure*, 33(3):321–343, 2000.
- [38] A. Yu. Ol’shanskii and M. V. Sapir. On F_k -like groups. *Algebra Logika*, 48(2):245–257, 284, 286–287, 2009.

OMER TAMUZ, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA, CA 91125.
 EMAIL: OMERTAMUZ@GMAIL.COM.

TIANYI ZHENG, UNIVERSITY OF CALIFORNIA, SAN DIEGO, LA JOLLA, CA 92093.
 EMAIL: TZHENG2@MATH.UCSB.EDU.